

Publication

Towards a better understanding of the generation of fructan structure diversity in plants: molecular and functional characterization of a sucrose:fructan 6-fructosyltransferase (6-SFT) cDNA from perennial ryegrass (*Lolium perenne*)

JournalArticle (Originalarbeit in einer wissenschaftlichen Zeitschrift)**ID** 1006548**Author(s)** Lasseur, B.; Lothier, J.; Wiemken, A.; Van Laere, A.; Morvan-Bertrand, A.; Van den Ende, W.; Prud'homme, M. P.**Author(s) at UniBasel** [Wiemken, Andres M.](#) ;**Year** 2011**Title** Towards a better understanding of the generation of fructan structure diversity in plants: molecular and functional characterization of a sucrose:fructan 6-fructosyltransferase (6-SFT) cDNA from perennial ryegrass (*Lolium perenne*)**Journal** Journal of Experimental Botany**Volume** 62**Number** 6**Pages / Article-Number** 1871-1885

The main storage compounds in *Lolium perenne* are fructans with prevailing beta(2-6) linkages. A cDNA library of *L. perenne* was screened using *Poa secunda* sucrose:fructan 6-fructosyltransferase (6-SFT) as a probe. A full-length Lp6-SFT clone was isolated as shown by heterologous expression in *Pichia pastoris*. High levels of Lp6-SFT transcription were found in the growth zone of elongating leaves and in mature leaf sheaths where fructans are synthesized. Upon fructan synthesis induction, Lp6-SFT transcription was high in mature leaf blades but with no concomitant accumulation of fructans. In vitro studies with the recombinant Lp6-SFT protein showed that both 1-kestotriose and 6G-kestotriose acted as fructosyl acceptors, producing 1- and 6-kestotetraose (bifurcose) and 6G,6-kestotetraose, respectively. Interestingly, bifurcose formation ceased and 6G,6-kestotetraose was formed instead, when recombinant fructan:fructan 6G-fructosyltransferase (6G-FFT) of *L. perenne* was introduced in the enzyme assay with sucrose and 1-kestotriose as substrates. The remarkable absence of bifurcose in *L. perenne* tissues might be explained by a higher affinity of 6G-FFT, as compared with 6-SFT, for 1-kestotriose, which is the first fructan formed. Surprisingly, recombinant 6-SFT from *Hordeum vulgare*, a plant devoid of fructans with internal glucosyl residues, also produced 6G,6-kestotetraose from sucrose and 6G-kestotriose. In the presence of recombinant *L. perenne* 6G-FFT, it produced 6G,6-kestotetraose from 1-kestotriose and sucrose, like *L. perenne* 6-SFT. Thus, we demonstrate that the two 6-SFTs have close catalytic properties and that the distinct fructans formed in *L. perenne* and *H. vulgare* can be explained by the presence of 6G-FFT activity in *L. perenne* and its absence in *H. vulgare*.

Publisher Oxford University Press**ISSN/ISBN** 0022-0957 ; 1460-2431**edoc-URL** <http://edoc.unibas.ch/46926/>**Full Text on edoc** No;**Digital Object Identifier DOI** 10.1093/jxb/erq388**ISI-Number** 000288553000012