

Publication

Crying babies, tired mothers: what do we know: a systematic review

JournalArticle (Originalarbeit in einer wissenschaftlichen Zeitschrift)

ID 1022868

Author(s) Kurth, E.; Kennedy, H. P.; Spichiger, E.; Hosli, I.; Zemp Stutz E.,

Author(s) at UniBasel Kurth, Elisabeth ; Zemp Stutz, Elisabeth ; Hösli-Krais, Irene M. ; Spichiger,

Elisabeth;

Year 2011

Title Crying babies, tired mothers: what do we know: a systematic review

Journal Midwifery

Volume 27

Number 2

Pages / Article-Number 187-194

Keywords Postpartum period, Infant crying, Colic, Postpartum fatigue

Mesh terms Chronobiology Disorders, prevention & control; Crying; Depression, Postpartum, prevention & control; Fatigue, psychology; Female; Humans; Infant Behavior; Infant Care, organization & administration; Infant, Newborn; Maternal Health Services, organization & administration; Maternal-Child Nursing, standards; Mother-Child Relations; Mothers; Postpartum Period, psychology

OBJECTIVE: to synthesise the evidence on the interconnectedness of infant crying and maternal tiredness in the postpartum period, both from quantitative as well as from qualitative studies. METHODS: a systematic review was conducted including studies in English, French and German published from 1980 to 2007. Studies were included in the systematic review if they had extractable data on infant crying as well as maternal tiredness in the period of 0-3 months post partum. Of 100 retrieved publications, 10 met these criteria. FINDINGS: evidence from this review indicated that the amount of infant crying during the first three months postpartum is associated with the experience of tiredness and fatigue in new mothers. Significant associations were found in five of six quantitative studies. The four identified qualitative studies describe how infant crying disrupts new mothers' circadian rhythms, reducing opportunities to rest and exacerbating tiredness. Incremental exhaustion diminished parents' ability to concentrate, raising the fear of harming their children, triggering depressive symptoms and burdening parent-child interaction. KEY CONCLUSIONS AND IMPLICATIONS FOR PRACTICE: if healthcare professionals are to address the prominent concerns of parents caring for a neonate, it is essential to review current care practices and tailor them to maternal and infant needs. A care strategy alleviating the burden of infant crying and maternal fatigue has the potential to strengthen family health from the earliest stage

Publisher Churchill Livingstone

ISSN/ISBN 0266-6138

edoc-URL http://edoc.unibas.ch/dok/A6002154

Full Text on edoc No;

Digital Object Identifier DOI 10.1016/j.midw.2009.05.012 **PubMed ID** http://www.ncbi.nlm.nih.gov/pubmed/19773102

ISI-Number WOS:000289071200013

Document type (ISI) Review