

## Publication

n-Alkane biosynthetic hydrogen isotope fractionation is not constant throughout the growing season in the riparian tree *Salix viminalis*

**JournalArticle (Originalarbeit in einer wissenschaftlichen Zeitschrift)****ID** 3342696**Author(s)** Newberry, Sarah L.; Kahmen, Ansgar; Dennis, Paul; Grant, Alastair**Author(s) at UniBasel** [Kahmen, Ansgar](#) ; [Newberry, Sarah](#) ;**Year** 2015**Title** n-Alkane biosynthetic hydrogen isotope fractionation is not constant throughout the growing season in the riparian tree *Salix viminalis***Journal** *Geochimica et cosmochimica acta***Volume** 165**Pages / Article-Number** 75-85

Compound-specific  $\delta^{2}\text{H}$  values of leaf wax n-alkanes have emerged as a potentially powerful paleohydrological proxy. Research suggests terrestrial plant n-alkane  $\delta^{2}\text{H}$  values are strongly correlated with meteoric water  $\delta^{2}\text{H}$  values, and may provide information on temperature, relative humidity, evaporation, and precipitation. This is based upon several assumptions, including that biosynthetic fractionation of n-alkanes during synthesis is constant within a single species. Here we present a multi-isotope study of the n-alkanes of riparian *Salix viminalis* growing in Norwich, UK. We measured n-alkane  $\delta^{2}\text{H}$ , leaf water  $\delta^{2}\text{H}$ , xylem water  $\delta^{2}\text{H}$ , and bulk foliar  $\delta^{13}\text{C}$  and evaluated the variability of n-alkane  $\delta^{2}\text{H}$  values and net biosynthetic fractionation ( $\epsilon_{\text{lw-wax}}$ ) over a whole growing season. *S. viminalis* n-alkane  $\delta^{2}\text{H}$  values decreased by 40% between the start of the growing season in April and the time when they stabilized in July. Variation in leaf and xylem water  $\delta^{2}\text{H}$  did not explain this variability.  $\epsilon_{\text{lw-wax}}$  varied from  $-116\%$  during leaf expansion in April to  $-156\%$  during the stable phase. This suggests that differential biosynthetic fractionation was responsible for the strong seasonal trends in *S. viminalis* n-alkane  $\delta^{2}\text{H}$  values. We suggest that variability in  $\epsilon_{\text{lw-wax}}$  is driven by seasonal differences in the carbohydrate source and thus the NADPH used in n-alkane biosynthesis, with stored carbohydrates utilized during spring and recent occurring growing season assimilates used later in the season. This is further supported by bulk foliar  $\delta^{13}\text{C}$  values, which are  $^{13}\text{C}$ -enriched during the period of leaf flush, relative to the end of the growing season. Our results challenge the assumption that biosynthetic fractionation is constant for a given species, and suggest that  $^{2}\text{H}$ -enriched stored assimilates are an important source for n-alkane biosynthesis early in the growing season. These findings have implications for the interpretation of sedimentary n-alkanes and call for a careful design of calibration studies using contemporary samples.

**Publisher** Elsevier Science**ISSN/ISBN** 0016-7037**edoc-URL** <http://edoc.unibas.ch/40144/>**Full Text on edoc** No;**Digital Object Identifier DOI** 10.1016/j.gca.2015.05.001**ISI-Number** WOS:000358427700006**Document type (ISI)** Article