

Publication

Do Swiss community pharmacists address the risk of sexually transmitted infections during a consultation on emergency contraception? A simulated patient study

JournalArticle (Originalarbeit in einer wissenschaftlichen Zeitschrift)

ID 4514422

Author(s) Haag, Melanie; Gudka, Sajni; Hersberger, Kurt E.; Arnet, Isabelle Author(s) at UniBasel Arnet, Isabelle; Hersberger, Kurt; Haag, Melanie;

Year 2019

Title Do Swiss community pharmacists address the risk of sexually transmitted infections during a consultation on emergency contraception? A simulated patient study

Journal European Journal of Contraception and Reproductive Health Care

Volume 24 Number 5

Pages / Article-Number 407-412

Keywords Community pharmacy; compliance with protocol; emergency contraception; sexually transmitted infection; simulated patient study

Objectives:; Since 2002, Swiss community pharmacists have dispensed emergency contraception (EC) as pharmacist-only medicine ideally using the official Swiss protocol. Our study aimed to determine pharmacists' resolution of an imaginary EC case, compliance with the protocol, and provision of information on the risk of sexually transmitted infections (STIs).; Methods:; We conducted a simulated patient study with 69 students who each visited a community pharmacy. The scenario started with the student requesting the 'morning after pill'. Current practice was assessed using an online evaluation form adapted from the Medication-Related Consultation Framework. Descriptive and statistical analyses were carried out.; Results:; All pharmacists correctly identified that the person needed EC. All pharmacists used an EC protocol and asked on average 10.9 (standard deviation 0.68) of 11 compulsory EC assessment questions. In total, 93% of pharmacists addressed EC counselling items and 56% addressed the risk of STIs, mainly by mentioning that condoms offered the best protection (76%).; Conclusions:; Community pharmacists correctly issued the EC, complied with the dispensing protocol and used their professional judgement to ensure optimal EC use. There is nevertheless room for improvement regarding pharmacists' STI counselling. Finally, the protocol's STI section could be enriched with specific information to guide counselling.

Publisher TAYLOR & FRANCIS LTD

ISSN/ISBN 1473-0782

edoc-URL https://edoc.unibas.ch/74489/

Full Text on edoc No;

Digital Object Identifier DOI 10.1080/13625187.2019.1661377 PubMed ID http://www.ncbi.nlm.nih.gov/pubmed/31526080

ISI-Number WOS:000487051200001

Document type (ISI) Article