

## Publication

### Protocol for MAAESTRO: Electronic Monitoring and Improvement of Adherence to Direct Oral Anticoagulant Treatment-A Randomized Crossover Study of an Educational and Reminder-Based Intervention in Ischemic Stroke Patients Under Polypharmacy

#### **Journalltem (Reviews, Editorials, Rezensionen, Urteilsanmerkungen etc. in einer wissenschaftlichen Zeitschrift)**

**ID** 4526314

**Author(s)** Polymeris, Alexandros A.; Albert, Valerie; Hersberger, Kurt E.; Engelter, Stefan T.; Schaedelin, Sabine; Arnet, Isabelle; Lyrer, Philippe A.

**Author(s) at UniBasel** [Arnet, Isabelle](#) ; [Polymeris, Alexandros](#) ; [Albert, Valerie](#) ; [Hersberger, Kurt](#) ; [Engelter, Stefan](#) ; [Lyrer, Philippe A.](#) ; [Schädelin, Sabine](#) ;

**Year** 2018

**Title** Protocol for MAAESTRO: Electronic Monitoring and Improvement of Adherence to Direct Oral Anticoagulant Treatment-A Randomized Crossover Study of an Educational and Reminder-Based Intervention in Ischemic Stroke Patients Under Polypharmacy

**Journal** Frontiers in Neurology

**Volume** 9

**Pages** 1134

**Keywords** adherence; adherence-improving intervention; direct oral anticoagulants; electronic monitoring; ischemic stroke; multi-compartment compliance aid; polypharmacy

Background: Non-adherence to direct oral anticoagulants (DOACs) remains a matter of concern, especially for patients with a recent stroke. However, data on electronically monitored adherence and adherence-improving interventions are scarce. Aims: We aim to use electronic monitoring in DOAC-treated stroke patients to (i) evaluate the effect of an educational, reminder-based adherence-improving intervention, (ii) investigate predictors of non-adherence, (iii) identify reliable self-report measures of adherence, and (iv) explore the association of non-adherence with clinical outcomes. Methods: Single-center, randomized, crossover, open-label study. Adherence to DOACs of polymedicated patients self-administering their medication will be monitored electronically throughout the 12-month-long study following hospitalization for ischemic stroke. After a 6-month observational phase, patients will receive pharmaceutical counseling with feedback on their intake history and be given a multi-compartment pillbox for the subsequent 6-month interventional phase. The pillbox will provide intake reminders either during the first or the last three interventional-phase months. Patients will be randomly allocated to reminders-first or reminders-last. Study outcomes: Primary: non-optimal timing adherence; Secondary: non-optimal taking adherence; timing adherence; taking adherence; self-reported adherence; clinical outcomes including ischemic and hemorrhagic events; patient-reported device usability and satisfaction. Sample size estimates: A sample of 130 patients provides 90% power to show a 20% improvement of the primary adherence outcome with intake reminders. Discussion: MAAESTRO will investigate various aspects of non-adherence and evaluate the effect of an adherence-improving intervention in DOAC-treated patients with a recent stroke using electronic monitoring. Clinical Trial Registration: ClinicalTrials.gov identifier: NCT03344146, Swiss National Clinical Trials Portal SNCTP000002410.

**Publisher** FRONTIERS MEDIA SA

**ISSN/ISBN** 1664-2295

**edoc-URL** <https://edoc.unibas.ch/74559/>

**Full Text on edoc** No;

**Digital Object Identifier DOI** 10.3389/fneur.2018.01134  
**PubMed ID** <http://www.ncbi.nlm.nih.gov/pubmed/30622509>  
**ISI-Number** WOS:000454099000001  
**Document type (ISI)** Article