

Publication

3C0Q: Uvde E175a

JournalArticle (Originalarbeit in einer wissenschaftlichen Zeitschrift)

ID 4531015

Author(s) Meulenbroek, E. M.; Paspaleva, K.; Thomassen, E. A. J.; Abrahams, J. P.; Goosen, N.; Pan-nu, N. S.

Author(s) at UniBasel [Abrahams, Jan Pieter](#) ;

Year 2008

Title 3C0Q: Uvde E175a

Journal Worldwide Protein Data Bank

Pages / Article-Number 3C0Q

Keywords Hydrolase

Mesh terms Science & TechnologyLife Sciences & BiomedicineBiochemistry & Molecular BiologyBio-chemistry & Molecular Biology

UV damage endonuclease is a DNA repair enzyme that can both recognize damage such as UV lesions and introduce a nick directly 5' to them. Recently, the crystal structure of the enzyme from *Thermus thermophilus* was solved. In the electron density map of this structure, unexplained density near the active site was observed at the tip of Lys229. Based on this finding, it was proposed that Lys229 is post-translationally modified. In this article, we give evidence that this modification is a carboxyl group. By combining activity assays and X-ray crystallography on several point mutants, we show that the carboxyl group assists in metal binding required for catalysis by donating negative charge to the metal-coordinating residue His231. Moreover, functional and structural analysis of the K229R mutant reveals that if His231 shifts away, an increased activity results on both damaged and undamaged DNA. Taken together, the results show that *T. thermophilus* ultraviolet damage endonuclease is carboxylated and the modified lysine is required for proper catalysis and preventing increased incision of undamaged DNA.

edoc-URL <https://edoc.unibas.ch/75944/>

Full Text on edoc No;

Digital Object Identifier DOI 10.2210/pdb3c0q/pdb

ISI-Number 2012140001768348

Document type (ISI) Data set