

Publication

1MO8: ATPase

JournalArticle (Originalarbeit in einer wissenschaftlichen Zeitschrift)

ID 4531050

Author(s) Hilge, M.; Siegal, G.; Vuister, G. W.; Guentert, P.; Gloor, S. M.; Abrahams, J. P.

Author(s) at UniBasel [Abrahams, Jan Pieter](#) ;

Year 2003

Title 1MO8: ATPase

Journal Worldwide Protein Data Bank

Pages / Article-Number 1MO8

Keywords Hydrolase

Mesh terms Science & TechnologyLife Sciences & BiomedicineBiochemistry & Molecular BiologyBiochemistry & Molecular Biology

The Na,K-ATPase hydrolyzes ATP to drive the coupled extrusion and uptake of Na⁺ and K⁺ ions across the plasma membrane. Here, we report two high-resolution NMR structures of the 213-residue nucleotide-binding domain of rat alpha1 Na,K-ATPase, determined in the absence and the presence of ATP. The nucleotide binds in the anti conformation and shows a relative paucity of interactions with the protein, reflecting the low-affinity ATP-binding state. Binding of ATP induces substantial conformational changes in the binding pocket and in residues located in the hinge region connecting the N- and P-domains. Structural comparison with the Ca-ATPase stabilized by the inhibitor thapsigargin, E2(TG), and the model of the H-ATPase in the E1 form suggests that the observed changes may trigger the series of events necessary for the release of the K⁺ ions and/or disengagement of the A-domain, leading to the eventual transfer of the gamma-phosphate group to the invariant Asp369.

edoc-URL <https://edoc.unibas.ch/75966/>

Full Text on edoc No;

Digital Object Identifier DOI 10.2210/pdb1mo8/pdb

ISI-Number 2012136001725619

Document type (ISI) Data set