

# Publication

Assessment of diabetes and prediabetes prevalence and predictors by HbA1c in a population from sub-Saharan Africa with a high proportion of anemia: a prospective cross-sectional study

## JournalArticle (Originalarbeit in einer wissenschaftlichen Zeitschrift)

## **ID** 4598467

**Author(s)** Hodel, Nikolai Carl; Hamad, Ali; Reither, Klaus; Mwangoka, Grace; Kasella, Irene; Praehauser, Claudia; Abdulla, Salim; Hatz, Christoph F. R.; Mayr, Michael

Author(s) at UniBasel Hodel, Nikolai ; Reither, Klaus ; Hatz, Christoph ;

#### Year 2020

**Title** Assessment of diabetes and prediabetes prevalence and predictors by HbA1c in a population from sub-Saharan Africa with a high proportion of anemia: a prospective cross-sectional study

Journal BMJ open diabetes research & care

Volume 8

#### Number 1

## Pages / Article-Number e000939

**Keywords** HbA1c; Tanzania; age; anemia; body mass index; diabetes; epidemiology; pre-diabetes; risk factors; sub-Sahara Africa

Epidemiological data about diabetes mellitus (DM) for sub-Saharan Africa (SSA) are scarce and the utility of glycated hemoglobin (HbA1c) to diagnose DM is uncertain in African populations with a high proportion of anemia.; In a cross-sectional study, age-adjusted prevalence rates and predictors for DM and pre-DM were prospectively assessed by HbA1c in a semirural walk-in population of Tanzania (n=992). Predictors for DM were calculated by logistic regression. Correlations between HbA1c, hemoglobin, and blood glucose levels were done by Pearson's correlation.; Overall, DM and pre-DM prevalence rates were 6.8% (95% CI 5.3 to 8.5) and 25% (95% CI 22.8 to 28.3), respectively. There was an increase in DM prevalence in patients 50-59 (14.9%; 95% CI 9.1 to 22.5), ≥60 years old (18.5%; 95% CI 12.2 to 26.2) and in patients with overweight (9.3%; 95% CI 5.9 to 13.7), obesity (10.9%; 95% CI 6.9 to 16) compared with patients 18-29 years old (2.2%; 95% CI 0.9 to 4.4) (p<0.001) and to normal-weight patients (3.6%; 95% CI 2.1 to 5.6) (p<0.01), respectively. Age (OR 1.08, 95% CI 1.05 to 1.12; p<0.001), body mass index (BMI) (OR 1.10, 95% CI 1.04 to 1.16; p<0.001), and acute infection (OR 3.46, 95% CI 1.02 to 10.8; p=0.038) were predictors for DM. Comparing patients with a BMI of 20 kg/m; 2; and a BMI of 35 kg/m; 2; , the relative risk for DM increases in average by 2.12-fold (range 1.91-2.24) across the age groups. Comparing patients 20 years old with patients 70 years old, the relative risk for DM increases in average 9.7-fold (range 8.9-10.4) across the BMI groups. Overall, 333 patients (36%) suffered from anemia. Pearson's correlation coefficients (r) between HbA1c and hemoglobin was -0.009 (p=0.779), and between HbA1c and fasting blood glucose and random blood glucose, it was 0.775 and 0.622, respectively (p<0.001).; We observed a high prevalence of DM and pre-DM, mainly triggered by increasing age and BMI, and provide evidence that HbA1c is suitable to assess DM also in populations of SSA with high proportions of anemia.; NCT03458338.

Publisher BMJ Publishing Group

ISSN/ISBN 2052-4897

edoc-URL https://edoc.unibas.ch/76860/

Full Text on edoc Available;

Digital Object Identifier DOI 10.1136/bmjdrc-2019-000939

PubMed ID http://www.ncbi.nlm.nih.gov/pubmed/32439737

ISI-Number MEDLINE:32439737 Document type (ISI) Journal Article

2