

Publication

Physical activity is favorably associated with arterial stiffness in patients with obesity and elevated metabolic risk

JournalArticle (Originalarbeit in einer wissenschaftlichen Zeitschrift)

ID 4606153

Author(s) Königstein, Karsten; Infanger, Denis; Klenk, Christopher; Carrard, Justin; Hinrichs, Timo; Schmidt-Trucksäss, Arno

Author(s) at UniBasel [Hinrichs, Timo](#) ; [Carrard, Justin](#) ;

Year 2020

Title Physical activity is favorably associated with arterial stiffness in patients with obesity and elevated metabolic risk

Journal The international journal of clinical practice

Volume 74

Number 9

Pages / Article-Number e13563

Mesh terms Adult; Blood Pressure; Body Composition; Cardiorespiratory Fitness, physiology; Exercise, physiology; Female; Humans; Male; Middle Aged; Motor Activity, physiology; Obesity, therapy; Pulse Wave Analysis; Vascular Stiffness

Physical activity and cardiorespiratory fitness (CRF) are relevant modifiers of cardiovascular risk. Their independent effects on arterial stiffness have not been assessed in people with obesity. This study aimed to assess the independent effects of light (LPA) and moderate-to-vigorous (MVPA) physical activity and CRF on Pulse wave velocity (PWV).; Brachial-ankle PWV (baPWV) was measured cross-sectionally in 55 subjects (43.0 ± 13.8 years; 66% women) with moderate cardiovascular risk. Body composition was assessed with bioelectrical impedance-analysis. Daily minutes of LPA and MVPA were measured by accelerometry and CRF (peak oxygen uptake [VO₂ peak]) with spiroergometry. Independent effects of LPA, MVPA, and VO₂ peak on baPWV were analyzed in an age-, sex-, body fat mass-, and blood pressure-adjusted ANOVA.; Every 10 minutes increase of daily MVPA was associated with a 2.8% (0.32m/s [-0.64 to 0.001 m/s], P = .05) reduction of baPWV, whereas LPA and VO₂ peak had only a little or no relevant effects on baPWV.; Higher MVPA is associated with lower composite arterial stiffness independent of CRF and the number of metabolic risk factors in patients with obesity and further metabolic risk factors. Thus, lifestyle interventions should aim for an increase in MVPA. BaPWV may improve the monitoring of favorable effects of MVPA, even if an improvement of VO₂ peak cannot be obtained.

Publisher Blackwell

ISSN/ISBN 1742-1241

edoc-URL <https://edoc.unibas.ch/79211/>

Full Text on edoc No;

Digital Object Identifier DOI 10.1111/ijcp.13563

PubMed ID <http://www.ncbi.nlm.nih.gov/pubmed/32478973>

ISI-Number WOS:000566922000024

Document type (ISI) Journal Article