

Publication

Explicitly accounting for needle sugar pool size crucial for predicting intraseasonal dynamics of needle carbohydrates $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$

JournalArticle (Originalarbeit in einer wissenschaftlichen Zeitschrift)**ID** 4646993**Author(s)** Leppä, Kersti; Tang, Yu; Ogée, Jérôme; Launiainen, Samuli; Kahmen, Ansgar; Kolari, Pasi; Sahlstedt, Elina; Saurer, Matthias; Schiestl-Aalto, Pauliina; Rinne-Garmston, Katja T.**Author(s) at UniBasel** [Kahmen, Ansgar](#) ;**Year** 2022**Title** Explicitly accounting for needle sugar pool size crucial for predicting intraseasonal dynamics of needle carbohydrates $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ **Journal** New phytologist**Pages / Article-Number** 18227**Keywords** Scots pine (*Pinus sylvestris*); boreal forest; carbon isotope; dynamic modeling; needle sugar; oxygen isotope; photosynthesis

We explore needle sugar isotopic compositions ($\delta^{18}\text{O}$ and $\delta^{13}\text{C}$) in boreal Scots pine (*Pinus sylvestris*) over two growing seasons. A leaf-level dynamic model driven by environmental conditions and based on current understanding of isotope fractionation processes was built to predict $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ of two hierarchical needle carbohydrate pools, accounting for the needle sugar pool size and the presence of an invariant pinitol pool. Model results agreed well with observed needle water $\delta^{18}\text{O}$, $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ of needle water-soluble carbohydrates (sugars + pinitol), and needle sugar $\delta^{13}\text{C}$ ($R^2 = 0.95, 0.84, 0.60, 0.73$, respectively). Relative humidity (RH) and intercellular to ambient CO_2 concentration ratio (C_i/C_a) were the dominant drivers of $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ variability, respectively. However, the variability of needle sugar $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ was reduced on diel and intra-seasonal timescales, compared to predictions based on instantaneous RH and C_i/C_a , due to the large needle sugar pool, which caused the signal formation period to vary seasonally from 2 d to more than 5 d. Furthermore, accounting for a temperature-sensitive biochemical ^{18}O -fractionation factor and mesophyll resistance in ^{13}C -discrimination were critical. Interpreting leaf-level isotopic signals requires understanding on time integration caused by mixing in the needle sugar pool.

Publisher Blackwell Science**ISSN/ISBN** 0028-646X ; 1469-8137**edoc-URL** <https://edoc.unibas.ch/89270/>**Full Text on edoc** No;**Digital Object Identifier DOI** 10.1111/nph.18227**PubMed ID** <http://www.ncbi.nlm.nih.gov/pubmed/35575976>**ISI-Number** WOS:000815022800001**Document type (ISI)** Article; Early Access