

## Publication

## Strengthening primary healthcare in Kosovo requires tailoring primary, secondary and tertiary prevention interventions and consideration of mental health

**JournalArticle (Originalarbeit in einer wissenschaftlichen Zeitschrift)**

ID 4651777

**Author(s)** Obas, K. A.; Bytyci-Katanolli, A.; Kwiatkowski, M.; Ramadani, Q.; Fota, N.; Jerliu, N.; Statovci, S.; Gerold, J.; Zahorka, M.; Probst-Hensch, N.**Author(s) at UniBasel** Obas, Katrina ; Bytyci Katanolli, Ariana ; Kwiatkowski, Marek ; Zahorka, Manfred ; Probst-Hensch, Nicole ;**Year** 2022**Title** Strengthening primary healthcare in Kosovo requires tailoring primary, secondary and tertiary prevention interventions and consideration of mental health**Journal** Front Public Health**Volume** 10**Pages / Article-Number** 794309**Keywords** Cross-Sectional Studies; Depression/diagnosis/epidemiology; \*Diabetes Mellitus/epidemiology/prevention & control; Humans; \*Hypertension; Kosovo/epidemiology; Mental Health; \*Noncommunicable Diseases; Primary Health Care; \*Pulmonary Disease, Chronic Obstructive/complications/epidemiology/prevention & control; Tertiary Prevention; Copd; depressive symptoms; diabetes; hypertension; prevention; public health; commercial or financial relationships that could be construed as a potential; conflict of interest.**Mesh terms** Cross-Sectional Studies; Depression, epidemiology; Diabetes Mellitus, prevention & control; Humans; Hypertension; Kosovo, epidemiology; Mental Health; Noncommunicable Diseases; Primary Health Care; Pulmonary Disease, Chronic Obstructive, prevention & control; Tertiary Prevention**Objectives:** Kosovo has the lowest life expectancy in the Balkans. Primary healthcare (PHC) plays an essential role in non-communicable disease (NCD) prevention. We described primary, secondary and tertiary prevention indicators in Kosovo and assessed their association with depressive symptoms. **Methods:** PHC users (n = 977) from the Kosovo NCD cohort baseline study were included. Depressive symptoms were assessed using the Depressive Anxiety Stress Scale-21. Cross-sectional associations between depressive symptoms and prevention indicators were quantified with mixed logistic regression models. **Results:** Poor nutrition (85%), physical inactivity (70%), obesity (53%), and smoking (21%) were common NCD risk factors. Many cases of hypertension (19%), diabetes (16%) and Chronic Obstructive Pulmonary Disease (COPD) (45%) remained undetected by a PHC professional. Uncontrolled hypertension (28%), diabetes (79%), and COPD (76%) were also common. Depressive symptoms were positively associated with physical inactivity (OR 1.02; 95% CI 1.00-1.05 per 1-point increase in DASS-21) and undetected COPD (OR 1.07; 95% CI 1.00-1.15), but inversely with undetected diabetes (OR 0.95; 95% CI 0.91-1.00). **Conclusions:** Continued attention and tailored modifications to primary, secondary and tertiary prevention in Kosovo are needed to narrow the Balkan health gap.**ISSN/ISBN** 2296-2565 (Electronic)2296-2565 (Linking)**URL** <https://doi.org/10.3389/fpubh.2022.794309>**edoc-URL** <https://edoc.unibas.ch/90724/>**Full Text on edoc** Available;**Digital Object Identifier DOI** 10.3389/fpubh.2022.794309**PubMed ID** <http://www.ncbi.nlm.nih.gov/pubmed/35480592>**ISI-Number** WOS:000792452200001

