

Publication

Nutrient use efficiency and arbuscular mycorrhizal root colonisation of winter wheat cultivars in different farming systems of the DOK long-term trial

JournalArticle (Originalarbeit in einer wissenschaftlichen Zeitschrift)

ID 487772

Author(s) Hildermann, Isabell; Messmer, Monika; Dubois, David; Boller, Thomas; Wiemken, Andres; Mäder, Paul

Author(s) at UniBasel [Boller, Thomas](#) ; [Wiemken, Andres M.](#) ;

Year 2010

Title Nutrient use efficiency and arbuscular mycorrhizal root colonisation of winter wheat cultivars in different farming systems of the DOK long-term trial

Journal Journal of the Science of Food and Agriculture

Volume 90

Number 12

Pages / Article-Number 2027-38

Keywords Organic farming, plant breeding, winter wheat, nutrient use efficiency, arbuscular mycorrhizal fungi

BACKGROUND: For organic farming, cultivars are required with high nutrient use efficiency under nutrient limited conditions. Arbuscular mycorrhizal fungi (AMF) are known to contribute to nutrient uptake under low input conditions. We compared nutrient use efficiency (NUE) of old and modern organically and conventionally bred cultivars in organic and conventional systems and assessed AMF-root colonisation (AMF-RC) in relation to nutrient concentrations. **RESULTS:** Cultivars and systems had a statistically significant effect on nitrogen (N) and phosphorus (P) concentrations and NUE parameters, whereas no genotype x environment interactions appeared. In contrast to N and P uptake, the NUE parameters were higher under organic than under conventional conditions. NUE for N increased with the year of release of cultivars. In the organic systems, the organically bred cultivars could not outperform the conventionally bred cultivars in grain yield and NUE parameters. AMF-RC was higher in the organic than in the conventional system, but did not differ among cultivars. **CONCLUSION:** Cultivars achieving high NUE in the organic systems were found among modern cultivars, irrespective of the breeding programme. Nutrient conditions during the breeding programme did not affect AMF-RC. No clear evidence was found that AMF symbiosis contributed more to nutrient concentrations under low input than under high input conditions.

Publisher Wiley Interscience

ISSN/ISBN 1097-0010

URL <http://onlinelibrary.wiley.com/doi/10.1002/jsfa.4048/abstract>

edoc-URL <http://edoc.unibas.ch/dok/A5842297>

Full Text on edoc No;

Digital Object Identifier DOI 10.1002/jsfa.4048

PubMed ID <http://www.ncbi.nlm.nih.gov/pubmed/20582996>

ISI-Number WOS:000281332000011

Document type (ISI) Journal Article